

<https://helda.helsinki.fi>

Moving out but keeping in touch : contacts between endoplasmic reticulum and lipid droplets

Salo, Veijo T.

2019-04

Salo , V T & Ikonen , E 2019 , ' Moving out but keeping in touch : contacts between
endoplasmic reticulum and lipid droplets ' , Current Opinion in Cell Biology , vol. 57 , pp.
64-70 . <https://doi.org/10.1016/j.ceb.2018.11.002>

<http://hdl.handle.net/10138/301334>

<https://doi.org/10.1016/j.ceb.2018.11.002>

cc_by_nc_nd

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.



Moving out but keeping in touch: contacts between endoplasmic reticulum and lipid droplets

Veijo T Salo^{1,2} and Elina Ikonen^{1,2}

The formation of neutral lipid filled and phospholipid monolayer engulfed lipid droplets (LDs) from the bilayer of the endoplasmic reticulum (ER) is an active area of investigation. This process harnesses the biophysical properties of the lipids involved and necessitates cooperation of protein machineries in both organelle membranes. Increasing evidence suggests that once formed, LDs keep close contact to the mother organelle and that this may be achieved via several, morphologically distinct and potentially functionally specialized connections. These may help LDs to dynamically respond to changes in lipid metabolic status sensed by the ER. In this review, we will discuss recent progress in understanding how LDs interact with the ER.

Addresses

¹ Faculty of Medicine, Dept. of Anatomy and HiLIFE, Univ. of Helsinki, Finland

² Minerva Foundation Institute for Medical Research, Helsinki, Finland

Current Opinion in Cell Biology 2019, **57**:64–70

This review comes from a themed issue on **Cell signalling**

Edited by **Wouter H Moolenaar** and **Tamás Balla**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 23rd November 2018

<https://doi.org/10.1016/j.ceb.2018.11.002>

0955-0674/© 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Lipid droplets (LDs) are ubiquitous intracellular storage organelles specialized in the storage of excess energy in the form of neutral lipids [1,2]. LDs are engaged in contacts with several organelles, but have the most intimate relationship with the ER [3,4]. The ER is not only the site of LD biogenesis, but may be a life-long partner of this organelle, enabling dynamic remodelling of LDs in response to cellular energy status and needs.

The neutral lipid core of LDs consists of triglycerides (TAG) and cholesterol esters, while the phospholipid monolayer is primarily composed of phosphatidylcholine (PC). The majority of both neutral lipid and PC synthesizing enzymes are not present on LDs [2], implying that contacts of LDs to the ER are needed to facilitate LD growth. The presence of a monolayer makes LDs unique

as cellular organelles. Its main function is to shield the hydrophobic core from the aqueous cytosolic environment, acting as surfactant stabilizing the droplet, and its biophysical properties confer specificity to the LD proteome [5]. Here, we review some of the recent advances in understanding LD forming sites in the ER as well as the morphology, stability and protein composition of ER–LD contacts.

LD forming sites in the ER

LDs originate from the ER that harbours the majority of enzymes required for neutral lipid synthesis. In mammalian cells, the terminal step of TAG synthesis is catalysed by DGAT1 and DGAT2 enzymes that reside in the ER, although DGAT2 and its yeast homologue can also be located at the LD monolayer [6–8]. In the current model of LD biogenesis, newly synthesized neutral lipids are initially relatively freely diffusing within the ER bilayer. In the absence of LDs, TAG synthesizing machinery appears dispersed around the ER [8,9], and it is not known whether these initial steps take place in specified subdomains. However, with rising local concentrations, neutral lipids have the tendency to aggregate, which might lead to formation of lens-like structures in the ER bilayer. Such lenses have been predicted by simulations [10] and also observed in yeast by electron microscopy [11]. Subsequently, the coalescence of neutral lipids buds out towards the cytosol, forming a nascent LD.

Several candidate proteins acting at the initial steps of LD formation have been identified, including members of the perilipin family, ACLS3, FITM2, lipin, seipin and Rab18. Perilipins may aid in LD budding by accumulating at ER regions containing high contents of neutral lipids or diacylglycerol (DAG) and may thus help to stabilize nascent forming LDs [12*,13–15]. ACSL3, a key acyltransferase in the generation of TAG in mammalian cells, is enriched at the sites of LD formation [16], producing possible localized metabolic hotspots for activated fatty acid generation [17*]. Lipin mediated DAG synthesis has been shown to be important for nascent LD formation in yeast [18], while FITM2 has been proposed to shuffle TAG to the forming LD [19,20] and control local DAG levels at LD forming sites [21*]. In yeast, FITM2 is present at LD forming sites and may facilitate LD budding towards the cytosol [11,21*]. Finally, seipin and Rab18 are enriched at ER–LD contact sites, with suggested functions therein (see below).

Furthermore, increasing evidence speaks for the importance of the local ER membrane lipid composition in the LD forming sites. The intrinsic curvature of phospholipids was found to regulate LD budding [21[•]]. Negatively curved lipids, such as DAG, facilitated LD embedding in the ER while positively curved lipids, such as lysolipids, facilitated budding out of the LD. In another study, conducted primarily with artificial LDs in model bilayer membranes, the size of LDs could be controlled by the phospholipid composition of the bilayer through effects on surface tension. Different phospholipids were found to have differential LD budding aptitudes, and decreasing surface tension (such as accumulation of lysolipids) favoured egress of neutral lipids, resulting in budding of small LDs [22^{••}]. Indeed, in simulations LD budding can occur spontaneously, that is without proteins [23]. Overall, these observations suggest that LD formation could be a process driven by lipids, with proteins playing important regulatory roles in controlling their local concentrations.

Further advances in the field necessitate more detailed information on the lipid composition of the LD forming ER subdomains. This is a challenging task but steps forward have recently been taken. In yeast, the concomitant depletion of seipin and Pex30, a peroxisomal biogenesis factor, prevented LD formation, with accumulation of TAG in the ER membrane [24^{••}]. Interestingly, this defect was largely rescued by modulation of the ER phospholipid composition. This prompted the authors to propose that in the absence of seipin and Pex30, LD formation was fully-dependent on the ER phospholipid composition. Remarkably, these findings also emphasize, together with other recent studies [25,26^{••}], the intimate relationship between LD and peroxisome biogenesis sites. While LDs and peroxisomes are structurally distinct, both form at ER subdomains and play important roles in lipid metabolism.

LDs as constituents of the ER network

The overall structure of the ER network, its dynamics and subdomain organization may contribute to LD formation and ER–LD junctions. Interestingly, key ER–LD contact proteins Rab18 and FITM2 have been shown to be important for ER tubular network morphology [27,28]. The high membrane curvature in tubules or tubule regulating proteins might somehow facilitate LD formation. In support of this, a number of proteins important for ER network formation and maintenance have been shown to effect LD biogenesis [29–33]. An interesting recent proposal is that ER shaping proteins could modulate the surface tension of the ER and thus effect LD budding [34].

The ER network is continuously reshaped by the cytoskeleton, and both acto-myosin and microtubule networks have been implicated in LD trafficking and

maintenance [1,35,36]. How ER–LD contacts are regulated during LD trafficking is not known, but in a recent study in Cos-7 cells ER–LD interactions were found not to be altered by disturbing the microtubule cytoskeleton [37]. However, there may be cell-type-specific differences, as the formation of ER–LD contacts was proposed to be kinesin-1-dependent in hepatocytes [38]. Given that cytoskeletal rearrangements may impose elastic stress on the membrane, they might also contribute to ER–LD contact stability, for example via alterations in ER or LD surface tension.

Studies in yeast have identified a subpopulation of LDs that form at nuclear-ER vacuole junctions and contain a specific subsets of LD proteins [39[•],40]. These LDs are important during nutritional stress and can be modulated by protein factors recently linked to seipin [40,41], suggesting that spatial coordination of ER–LD contact sites may play a role in LD diversification.

Morphological features of ER–LD contacts


ER–LD contacts with variable morphology have been reported. These include lipidic bridges, where the LD monolayer appears to be directly continuous with the cytosolic leaflet of the ER bilayer [4]. Such discrete, relatively well-defined and small-sized contacts are visible in electron microscopic images in numerous cell types [7,42[•],43,44,45[•],46–51] (see also Figure 1). This also seems to hold for the recently characterized intranuclear LDs [52,53]. These connections often appear to have a finger-like protrusion extending from the ER. More extensive ER–LD contacts, where the LD appears to lie like an egg in a cup, have also been observed (for example in [8,54,55] and Figure 1). Finally, proteinaceous tethers between the ER and LDs have been noted [42[•]]. While it is likely that contacts with variable appearance are functionally distinct, direct evidence for this is still lacking.

Extent and stability of ER–LD contacts

Mounting evidence suggests that the formed LDs remain associated with the ER [8,45[•],46,54] and in fact, there is little evidence for total dissociation of LDs from the ER. By quantitative 3D-EM in human A431 cells, we found all LDs to be associated with ER membranes both during early and later stages of LD biogenesis [45[•]], and LDs without ER connections were only observed in seipin-deficient cells. High resolution live cell imaging of Cos-7 cells also indicated that virtually all LDs are in contact with ER membranes [37]. However, mature LDs in HeLa cells may not be fully associated with the ER, based on FRAP experiments [47]. So far, machineries facilitating ER–LD detachment have not been identified.

How long-lasting individual ER–LD contacts are, is at present not well understood. Studies in *Drosophila* cells suggest that they may be dynamic and form in response to

Figure 1



Examples of ER-LD contacts in human cells. **(a)** Live cell Airyscan microscopy of a human A431 cell grown in complete medium and treated with oleic acid for 2 h, expressing BFP-KDEL (ER, cyan), endogenously tagged seipin (magenta) and stained with a lipid droplet dye (LD540, yellow). Scale bars 5 and 1 μm . **(b)** 3D models of LDs and surrounding ER profiles shown on top of a block-face image from a serial block-face scanning EM dataset of a A431 cell treated with oleic acid for 20 h. Yellow indicates ER, brown LDs and red membrane proximities between ER and LDs. Scale bar 1 μm . **(c)**. Exemplary snapshots of ER-LD contacts from EM tomogram slices. Scale bars 200 nm. For detailed methodology, see Ref. [45].

alterations in the properties of the phospholipid monolayer [7,56]. In these studies, bridges were concluded to be important for the trafficking of lipogenic enzymes from the ER to a subset of LDs to allow subsequent LD expansion [7] and this trafficking was dependent on COPI machinery [56]. As COPI machinery can modulate the

LD surface tension by decreasing the monolayer PC levels (through budding out of small nano-LDs), the authors suggested that such decrease in surface tension may facilitate the bridge formation [5,56]. In addition, the LD monolayer phospholipid composition may be modulated by resident LPCAT enzymes [57] and localized

Table 1			
Proteins proposed to function at ER-LD contacts			
Name	Proposed function at ER-LD contacts	Localization ^a	References
Seipin	Regulating phosphatidic acid levels, functioning as a scaffold for lipogenic enzymes, as a diffusion barrier, or as a structural component facilitating LD growth	ER, enriched at ER-LD contacts	[42*,45*,60,61,64,65]
FITM2	Shuffling TAG to forming LDs, decreasing DAG at LD-forming sites to allow for LD emergence, regulating LD-budding to cytosol	ER, in yeast localised to ER-LD contacts	[11,19,20,21*]
Rab18	Inducing apposition of LDs to ER, interacts with NRZ-SNARE proteins at ER-LD interface, COPI-TRAPP1 complex also regulates its LD localisation	ER and LD monolayer	[68, 70*,72]
DGAT2-FATP1	Forms a complex to facilitate LD expansion	ER (FATP1) and LD (DGAT2) in <i>C. elegans</i>	[74]
ORP2-VAPA	May regulate hydrolysis, ORP2 depletion increases LD proximity to ER	ER (VAPA), multiple localizations including LD (ORP2)	[75,76]
VMP1	May modulate SERCA activity, knockdown increases LD proximity to ER	ER, overexpressed protein localized at multiple ER contacts, including ER-LD contacts.	[50,77]
ICE2	Channelling DAG from LDs to ER	ER and ER-LD contacts in yeast.	[78]

^a Localization in multiple cell types unless specified.

lipid transport. Interestingly, recent observations suggest that VPS13 proteins localize at ER contact sites, including those between ER and LDs, and may act as phospholipid transporters [58].

ER-LD contact proteins

The molecular machineries bridging the ER to LDs are becoming unravelled, with an increasing number of proteins localized and proposed to function at ER-LD contacts (Table 1).

The most studied of these proteins is seipin/BSCL2, a homo-oligomeric ER transmembrane protein of unknown function, mutated in severe lipodystrophy [59–61]. Seipin has been localised at ER-LD contacts in both yeast and human cells [42*,45*,60,61] and it plays an important role in LD biogenesis. In the absence of seipin, cells generate supersized and tiny, growth abortive LDs and exhibit functional defects in ER-LD contacts [42*,44,45*,52]. It is unclear whether the LD abnormalities arise from defective LD biogenesis or if seipin plays a role also in ER-LD contact maintenance. However, both nascent and more mature LDs harbour seipin at the ER-LD contact site and by live cell imaging these associations appear stable at least in minutes time-scale [42*,45*].

At ER-LD contacts seipin may regulate the local lipid environment, particularly phosphatidic acid levels [62–64], or act as a diffusion barrier between these organelles [44]. Seipin has also been suggested to act as a scaffold for lipid processing enzymes [65], or as a structural component of the ER-LD contacts facilitating LD growth [42*,45*]. Recently, the structures of the ER luminal regions of human and *drosophila* seipin oligomers have been solved using cryo-electron microscopy [66*,67*]. These structures indicate a ring-shaped oligomer of ~15–20 nm in diameter, with 11 (human) and 12 (*drosophila*) monomers. The luminal region harbours a β sandwich fold with structural similarity to lipid binding proteins, such as NPC2 [66*,67*]. These data suggest that seipin may function as a lipid transporter, but the putative lipid substrate(s) *in vivo* await further characterization.

Another protein implicated in ER-LD contacts is Rab18. Rab18 is localized to the ER and LDs [27,68,69,70*], and its LD localization is regulated by multiple factors including cellular metabolic status [71], COP1-TRAPP complex [72] and Rab3GAP1/2 [70*]. Rab18 has been linked to both lipolysis and lipogenesis [71], and its overexpression can induce close apposition of ER and LDs [68]. Recently, Rab18 was shown to bring together ER and LDs through an interaction with the NRZ complex and associated SNARE proteins, and depletion of this machinery resulted in reduced LD growth and maturation [70*]. These data suggest that the Rab18-NRZ complex might function as a tethering protein between

ER and LDs, although this function is apparently not important for LD growth in all cell types [70*,73].

Concluding remarks

In the past couple of years, we have witnessed substantial progress in understanding the connections between the ER and the lipid storage organelle it can give rise to, the LD. This has prompted a number of exciting avenues for further research. For instance, it will be interesting to dissect the functions of ER-LD contact proteins beyond their roles in initial LD formation, including LD maintenance and lipolysis. Furthermore, combining high resolution techniques, especially EM, with dynamic measurements and spatially resolved lipidomics will be instrumental for understanding the functional specifications of the various kinds of ER-LD contacts identified.

Conflict of interest statement

Nothing declared.

Acknowledgements

We thank HiLIFE Light and Electron microscopy platforms for imaging support, and Leena Karhinen, Ilya Belevich and Helena Vihinen for providing images. This study was supported by grants from the Academy of Finland (282192, 307415, 312491), Sigrid Juselius Foundation, University of Helsinki (HiLIFE Fellowship), Finnish Medical Foundation, Paulo Foundation, Alfred Kordelin Foundation, Maud Kuistila Foundation, Biomedicum Helsinki Foundation and Emil Aaltonen Foundation.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Welte MA, Gould AP: **Lipid droplet functions beyond energy storage.** *Biochim Biophys Acta - Mol Cell Biol Lipids* 2017, **1862**:1260–1272.
 2. Henne WM, Reese ML, Goodman JM: **The assembly of lipid droplets and their roles in challenged cells.** *EMBO J* 2018, **37**: e98947.
 3. Ohsaki Y, Soltysik K, Fujimoto T: **The lipid droplet and the endoplasmic reticulum.** *Organelle Contact Sites*. Singapore: Springer; 2017, 111–120.
 4. Schuldiner M, Bohnert M: **A different kind of love – lipid droplet contact sites.** *Biochim Biophys Acta - Mol Cell Biol Lipids* 2017, **1862**:1188–1196.
 5. Thiam AR, Farese RV, Walther TC: **The biophysics and cell biology of lipid droplets.** *Nat Rev Mol Cell Biol* 2013, **14**:775–786.
 6. Kuerschner L, Moessinger C, Thiele C: **Imaging of lipid biosynthesis: how a neutral lipid enters lipid droplets.** *Traffic* 2008, **9**:338–352.
 7. Wilfling F, Wang H, Haas JT, Krahmer N, Gould TJ, Uchida A, Cheng J-X, Graham M, Christiano R, Fröhlich F *et al.*: **Triacylglycerol synthesis enzymes mediate lipid droplet growth by relocating from the ER to lipid droplets.** *Dev Cell* 2013, **24**:384–399.
 8. Jacquier N, Choudhary V, Mari M, Toulmay A, Reggiori F, Schneider R: **Lipid droplets are functionally connected to the endoplasmic reticulum in *Saccharomyces cerevisiae*.** *J Cell Sci* 2011, **124**:2424–2437.
 9. Stone SJ, Levin MC, Zhou P, Han J, Walther TC, Farese RV: **The endoplasmic reticulum enzyme DGAT2 is found in mitochondria-associated membranes and has a**

mitochondrial targeting signal that promotes its association with mitochondria. *J Biol Chem* 2009, **284**:5352-5361.

10. Khandelia H, Duelund L, Pakkanen KI, Ipsen JH: **Triglyceride blisters in lipid bilayers: implications for lipid droplet biogenesis and the mobile lipid signal in cancer cell membranes.** *PLoS One* 2010, **5**:1-8.
 11. Choudhary V, Ojha N, Golden A, Prinz WA: **A conserved family of proteins facilitates nascent lipid droplet budding from the ER.** *J Cell Biol* 2015, **211**:261-271.
 12. Gao Q, Binns DD, Kinch LN, Grishin NV, Ortiz N, Chen X, Goodman JM: **Pet10p is a yeast perilipin that stabilizes lipid droplets and promotes their assembly.** *J Cell Biol* 2017, **216**:3199-3217.
- Pet10 was found to function as a yeast perilipin protein. It was highly abundant in TAG containing LDs and stabilized them, preventing LD fusion and facilitating LD biogenesis. In the absence of Pet10, TAG accumulation was reduced, but this was primarily due to defective TAG synthesis rather than increased lipolysis. Pet10 was found to functionally co-operate with seipin and FITM2 in LD formation.
13. Jacquier N, Mishra S, Choudhary V, Schneiter R: **Expression of oleosin and perilipins in yeast promotes formation of lipid droplets from the endoplasmic reticulum.** *J Cell Sci* 2013, **126**:5198-5209.
 14. Bulankina AV, Deggerich A, Wenzel D, Mutenda K, Wittmann JG, Rudolph MG, Burger KNJ, Höning S: **TIP47 functions in the biogenesis of lipid droplets.** *J Cell Biol* 2009, **185**:641-655.
 15. Skinner JR, Shew TM, Schwartz DM, Tzekov A, Lepus CM, Abumrad NA, Wolins NE: **Diacylglycerol enrichment of endoplasmic reticulum or lipid droplets recruits perilipin 3/ TIP47 during lipid storage and mobilization.** *J Biol Chem* 2009, **284**:30941-30948.
 16. Kassan A, Herms A, Fernández-Vidal A, Bosch M, Schieber NL, Reddy BJN, Fajardo A, Gelabert-Baldrich M, Tebar F, Enrich C *et al.*: **Acyl-CoA synthetase 3 promotes lipid droplet biogenesis in ER microdomains.** *J Cell Biol* 2013, **203**:985-1001.
 17. Poppelreuther M, Sander S, Minden F, Dietz MS, Exner T, Du C, Zhang I, Ehehalt F, Knüppel L, Domschke S *et al.*: **The metabolic capacity of lipid droplet localized acyl-CoA synthetase 3 is not sufficient to support local triglyceride synthesis independent of the endoplasmic reticulum in A431 cells.** *Biochim Biophys Acta - Mol Cell Biol Lipids* 2018, **1863**:614-624.
- This study reported that ACSL3 is the dominant acyl-CoA generating enzyme in human A431 cells, present in ER membranes and enriched in LDs. The authors also found that the metabolic activity of ACSL3 on LDs cannot explain the growth and expansion of LDs in these cells, implying that ongoing neutral lipid transport is required for LD expansion.
18. Adeyo O, Horn PJ, Lee S, Binns DD, Chandrasas A, Chapman KD, Goodman JM: **The yeast lipid orthologue Pah1p is important for biogenesis of lipid droplets.** *J Cell Biol* 2011, **192**:1043-1055.
 19. Gross DA, Zhan C, Silver DL: **Direct binding of triglyceride to fat storage-inducing transmembrane proteins 1 and 2 is important for lipid droplet formation.** *Proc Natl Acad Sci U S A* 2011, **108**:19581-19586.
 20. Kadereit B, Kumar P, Wang W-J, Miranda D, Snapp EL, Severina N, Torregroza I, Evans T, Silver DL: **Evolutionarily conserved gene family important for fat storage.** *Proc Natl Acad Sci U S A* 2008, **105**:94-99.
 21. Choudhary V, Golani G, Joshi AS, Cottier S, Schneiter R, Prinz WA, Kozlov MM: **Architecture of lipid droplets in endoplasmic reticulum is determined by phospholipid intrinsic curvature.** *Curr Biol* 2018, **28**:915-926.
- In silico* studies indicated that the intrinsic curvature of ER phospholipids influences ER-LD contact sites. In yeast, manipulating phospholipids affected LD emergence from the ER: lipids with high negative curvature, such as DAG, favoured LD embedding in the ER, while lipids with high positive curvature, such as lysolipids, favoured LD emergence. Accumulation of DAG at LD forming sites was suggested to be the culprit of LD budding defects observed in FITM2 knockout cells.
22. Ben M'barek K, Ajajji D, Chorlay A, Vanni S, Forêt L, Thiam AR: **ER membrane phospholipids and surface tension control cellular lipid droplet formation.** *Dev Cell* 2017, **41**:591-604.

Utilizing primarily artificial LDs and model membranes, the authors demonstrated that the bilayer phospholipid composition and surface tension are key parameters of LD budding. Investigation of individual phospholipid species revealed that low surface tension inducing lipids (such as lysolipids) induced LD budding and favoured formation of small LDs. Thus, altering ER phospholipid content or tension enables controlled LD formation.

23. Deslandes F, Thiam AR, Forêt L: **Lipid droplets can spontaneously bud off from a symmetric bilayer.** *Biophys J* 2017, **113**:15-18.
 24. Wang S, Idrissi FZ, Hermansson M, Grippa A, Ejsing CS, Carvalho P: **Seipin and the membrane-shaping protein Pex30 cooperate in organelle budding from the endoplasmic reticulum.** *Nat Commun* 2018, **9**:2939.
- Together with the work of Joshi *et al.***, an intimate link between LD and peroxisome biogenesis was reported. Seipin and Pex30 functionally interact in yeast, and the absence of both proteins resulted in major LD and peroxisome budding defects, which could be partially rescued by remodelling cellular phospholipid composition. The authors suggested that these proteins organize ER subdomains with a potentially unique lipid composition.
25. Schrüel B, Kopito RR: **Peroxisome-dependent targeting of a lipid droplet-destined membrane protein to ER subdomains.** *Nat Cell Biol* 2016, **18**:740-751.
 26. Joshi AS, Nebenfuhr B, Choudhary V, Satpute-Krishnan P, Levine TP, Golden A, Prinz WA: **Lipid droplet and peroxisome biogenesis occur at the same ER subdomains.** *Nat Commun* 2018, **9**:2940.
- Together with the work of Wang *et al.***, an intimate link between LD and peroxisome biogenesis was reported. Pex30 and seipin co-localized at LD forming sites, and mature LDs typically remained associated with Pex30 domains. MCTP2 was identified as a putative functional homologue of Pex30 in higher eukaryotes, and its depletion resulted in reduced LD generation in mammalian cells and in *C. elegans*.
27. Gerondopoulos A, Bastos RN, Yoshimura SI, Anderson R, Carpanini S, Aligianis I, Handley MT, Barr FA: **Rab18 and a Rab18 GEF complex are required for normal ER structure.** *J Cell Biol* 2014, **205**:707-720.
 28. Hayes MJ, Choudhary V, Ojha N, Shin JJ, Han G-S, Carman GM, Loewen CJ, Prinz WA, Levine TP: **Fat storage-inducing transmembrane (FIT or FITM) proteins are related to lipid phosphatase/phosphotransferase enzymes.** *Microb Cell* 2018, **5**:88-103.
 29. Klemm RW, Norton JP, Cole RA, Li CS, Park SH, Crane MM, Li L, Jin D, Boye-Doe A, Liu TY *et al.*: **A conserved role for atlastin GTPases in regulating lipid droplet size.** *Cell Rep* 2013, **3**:1465-1475.
 30. Renvoisé B, Malone B, Falgairolle M, Munasinghe J, Stadler J, Sibilla C, Park SH, Blackstone C: **Reep1 null mice reveal a converging role for hereditary spastic paraplegia proteins in lipid droplet regulation.** *Hum Mol Genet* 2016, **25**:5111-5125.
 31. Falk J, Rohde M, Bekhte MM, Neugebauer S, Hemmerich P, Kiehnopf M, Deufel T, Hübner CA, Beetz C: **Functional mutation analysis provides evidence for a role of REEP1 in lipid droplet biology.** *Hum Mutat* 2014, **35**:497-504.
 32. Xiang R, Fan L-L, Huang H, Chen Y-Q, He W, Guo S, Li J-J, Jin J-Y, Du R, Yan R *et al.*: **Increased RTN3 leads to obesity and hypertriglyceridemia by interacting with HSPA5.** *Circulation* 2018, **138**:1828-1838.
 33. Papadopoulos C, Orso G, Mancuso G, Herholz M, Gumeni S, Tadepalle N, Jüngst C, Tzschichholz A, Schauss A, Höning S *et al.*: **Spastin binds to lipid droplets and affects lipid metabolism.** *PLoS Genet* 2015, **11**:e1005149.
 34. Chorlay A, Thiam AR: **An asymmetry in monolayer tension regulates lipid droplet budding direction.** *Biophys J* 2018, **114**:631-640.
 35. Herms A, Bosch M, Reddy BJN, Schieber NL, Fajardo A, Rupérez C, Fernández-Vidal A, Ferguson C, Rentero C, Tebar F *et al.*: **AMPK activation promotes lipid droplet dispersion on detyrosinated microtubules to increase mitochondrial fatty acid oxidation.** *Nat Commun* 2015, **6**:7176.

36. Pfisterer SG, Gateva G, Horvath P, Pirhonen J, Salo VT, Karhinen L, Varjosalo M, Ryhänen SJ, Lappalainen P, Ikonen E: **Role for formin-like 1-dependent acto-myosin assembly in lipid droplet dynamics and lipid storage.** *Nat Commun* 2017, **8**:14858.
 37. Valm AM, Cohen S, Legant WR, Melunis J, Hershberg U, Wait E, Cohen AR, Davidson MW, Betzig E, Lippincott-Schwartz J: **Applying systems-level spectral imaging and analysis to reveal the organelle interactome.** *Nature* 2017, **546**:162-167.
 38. Rai P, Kumar M, Sharma G, Barak P, Das S, Kamat SS, Mallik R: **Kinesin-dependent mechanism for controlling triglyceride secretion from the liver.** *Proc Natl Acad Sci U S A* 2017, **114**:12958-12963.
 39. Hariri H, Rogers S, Ugrankar R, Liu YL, Feathers JR, Henne WM: **Lipid droplet biogenesis is spatially coordinated at ER-vacuole contacts under nutritional stress.** *EMBO Rep* 2018, **19**:57-72.
- This study found that during nutritional stress, yeast cells expanded their ER-vacuole contact sites (NVJ). LDs were often formed at sites near the NVJ and were required for NVJ expansion. Overexpression of NVJ-tether Mdm1 increased LD production. Therefore, spatial co-ordination of LD biogenesis sites within cells was concluded to be important during nutritional challenge.
40. Eisenberg-Bord M, Mari M, Weill U, Rosenfeld-Gur E, Moldavski O, Castro IG, Soni KG, Harpaz N, Levine TP, Futerman AH *et al.*: **Identification of seipin-linked factors that act as determinants of a lipid droplet subpopulation.** *J Cell Biol* 2018, **217**:269-282.
 41. Teixeira V, Johnsen L, Martínez-Montañés F, Grippa A, Buxó L, Idrissi FZ, Ejlsing CS, Carvalho P: **Regulation of lipid droplets by metabolically controlled Ldo isoforms.** *J Cell Biol* 2018, **217**:127-138.
 42. Wang H, Becuwe M, Housden BE, Chitraju C, Porras AJ, Graham MM, Liu XN, Thiam AR, Savage DB, Agarwal AK *et al.*: **Seipin is required for converting nascent to mature lipid droplets.** *Elife* 2016, **5**:133-150.
- An ER-to-LD marker peptide, LiveDrop, was developed, allowing early detection of nascent LDs. Using this tool and EM, the authors showed that in the absence of seipin, mammalian cells accumulated numerous tiny LDs that failed to grow. Endogenously tagged seipin was localized to ER-LD contacts, and the authors proposed that seipin helps nascent LDs to mature.
43. Wolinski H, Kolb D, Hermann S, Koning RI, Kohlwein SD: **A role for seipin in lipid droplet dynamics and inheritance in yeast.** *J Cell Sci* 2011, **124**:3894-3904.
 44. Grippa A, Buxó L, Mora G, Funaya C, Idrissi F-Z, Mancuso F, Gomez R, Muntanya J, Sapidó E, Carvalho P: **The seipin complex Fld1/Ldb16 stabilizes ER-lipid droplet contact sites.** *J Cell Biol* 2015, **211**:829-844.
 45. Salo VT, Belevich I, Li S, Karhinen L, Vihinen H, Vigouroux C, Magré J, Thiele C, Hölttä-Vuori M, Jokitalo E *et al.*: **Seipin regulates ER-lipid droplet contacts and cargo delivery.** *EMBO J* 2016, **24**:2699-2716.
- The authors used serial block-face scanning EM to study ER-LD contacts in human cells and found that all LDs were connected to the ER. Upon knocking out seipin, cells accumulated numerous tiny LDs, some of which were completely detached from the ER, and transport of lipid and protein constituents from ER to LDs became defective. Endogenously tagged seipin was localized to ER-LD contacts, and seipin was proposed to regulate ER-LD cargo delivery.
46. Ohsaki Y, Cheng J, Suzuki M, Fujita A, Fujimoto T: **Lipid droplets are arrested in the ER membrane by tight binding of lipidated apolipoprotein B-100.** *J Cell Sci* 2008, **121**:2415-2422.
 47. Kimura H, Arasaki K, Ohsaki Y, Fujimoto T, Ohtomo T, Yamada J, Tagaya M: **Syntaxin 17 promotes lipid droplet formation by regulating the distribution of acyl-CoA synthetase 3.** *J Lipid Res* 2018, **59**:805-819.
 48. Brocard L, Immel F, Coulon D, Esnay N, Tuphile K, Pascal S, Claverol S, Fouillen L, Bessoule J-J, Bréhélin C: **Proteomic analysis of lipid droplets from arabidopsis aging leaves brings new insight into their biogenesis and functions.** *Front Plant Sci* 2017, **8**.
 49. Wolf SG, Mutsafi Y, Dadosh T, Ilani T, Lansky Z, Horowitz B, Rubin S, Elbaum M, Fass D: **3D visualization of mitochondrial solid-phase calcium stores in whole cells.** *Elife* 2017, **6**:1-18.
 50. Zhao YG, Chen Y, Miao G, Zhao H, Qu W, Li D, Wang Z, Liu N, Li L, Chen S *et al.*: **The ER-localized transmembrane protein EPG-3/VMP1 regulates SERCA activity to control ER-isolation membrane contacts for autophagosome formation.** *Mol Cell* 2017, **67**:974-989.
 51. Wu H, Carvalho P, Voeltz GK: **Here, there, and everywhere: the importance of ER membrane contact sites.** *Science* 2018, **361**.
 52. Romanauska A, Köhler A: **The inner nuclear membrane is a metabolically active territory that generates nuclear lipid droplets.** *Cell* 2018, **174**:700-715.
 53. Ohsaki Y, Kawai T, Yoshikawa Y, Cheng J, Jokitalo E, Fujimoto T: **PML isoform II plays a critical role in nuclear lipid droplet formation.** *J Cell Biol* 2016, **212**:29-38.
 54. Mishra S, Khaddaj R, Cottier S, Stradalova V, Jacob C, Schneider R: **Mature lipid droplets are accessible to ER luminal proteins.** *J Cell Sci* 2016, **129**:3803-3815.
 55. Robenek H, Hofnagel O, Buers I, Robenek MJ, Troyer D, Severs NJ: **Adipophilin-enriched domains in the ER membrane are sites of lipid droplet biogenesis.** *J Cell Sci* 2006, **119**:4215-4224.
 56. Wilfling F, Thiam AR, Olarte M-J, Wang J, Beck R, Gould TJ, Allgeyer ES, Pincet F, Bewersdorf J, Farese RV *et al.*: **Arf1/COP1 machinery acts directly on lipid droplets and enables their connection to the ER for protein targeting.** *Elife* 2014, **3**:e01607.
 57. Moessinger C, Kuerschner L, Spandl J, Shevchenko A, Thiele C: **Human lysophosphatidylcholine acyltransferases 1 and 2 are located in lipid droplets where they catalyze the formation of phosphatidylcholine.** *J Biol Chem* 2011, **286**:21330-21339.
 58. Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, Wheeler H, Reinisch KM, De Camilli P: **VPS13A and VPS13C are lipid transport proteins differentially localized at ER contact sites.** *J Cell Biol* 2018, **217**:3625-3639.
 59. Magré J, Delépine M, Khallouf E, Gedde-Dahl T, Van Maldergem L, Sobel E, Papp J, Meier M, Mégarbané A, Bachy A *et al.*: **Identification of the gene altered in Berardinelli-Seip congenital lipodystrophy on chromosome 11q.** *Nat Genet* 2001, **28**:365-370.
 60. Szymanski KM, Binns D, Bartz R, Grishin NV, Li W, Agarwal AK, Garg A, Anderson RGW, Goodman JM: **The lipodystrophy protein seipin is found at endoplasmic reticulum lipid droplet junctions and is important for droplet morphology.** *Proc Natl Acad Sci U S A* 2007, **104**:20890-20895.
 61. Fei W, Shui G, Gaeta B, Du X, Kuerschner L, Li P, Brown J, Wenk MR, Parton RG, Yang H: **Fld1p, a functional homologue of human seipin, regulates the size of lipid droplets in yeast.** *J Cell Biol* 2008, **180**:473-482.
 62. Wolinski H, Hofbauer HF, Hellauer K, Cristobal-Sarramian A, Kolb D, Radulovic M, Knittelfelder OL, Rechberger GN, Kohlwein SD: **Seipin is involved in the regulation of phosphatidic acid metabolism at a subdomain of the nuclear envelope in yeast.** *Biochim Biophys Acta* 2015, **1851**:1450-1464.
 63. Han S, Binns DD, Chang Y-F, Goodman JM: **Dissecting seipin function: the localized accumulation of phosphatidic acid at ER/LD junctions in the absence of seipin is suppressed by Sei1pΔNterm only in combination with Ldb16p.** *BMC Cell Biol* 2015, **16**:29.
 64. Pagac M, Cooper DE, Qi Y, Lukmantara IE, Mak HY, Wu Z, Tian Y, Liu Z, Lei M, Du X *et al.*: **SEIPIN regulates lipid droplet expansion and adipocyte development by modulating the activity of glycerol-3-phosphate acyltransferase.** *Cell Rep* 2016, **17**:1546-1559.
 65. Talukder MMU, Sim MFM, O'Rahilly S, Edwardson JM, Rochford JJ: **Seipin oligomers can interact directly with AGPAT2 and lipin 1, physically scaffolding critical regulators of adipogenesis.** *Mol Metab* 2015, **4**:199-209.

66. Yan R, Qian H, Lukmantara I, Gao M, Du X, Yan N, Yang H: **Human • SEIPIN binds anionic phospholipids.** *Dev Cell* 2018, **47**:1-9.
Using cryo-EM, the authors report the structure of the luminal domain of human seipin, which exists as a ring-shaped undecamer. In vitro experiments indicate that seipin can bind anionic phospholipids.
67. Sui X, Arlt H, Brook K, Lai ZW, DiMaio F, Marks D, Liao M, Farese R, Walther T: **Cryo-electron microscopy structure of the lipid droplet-formation protein seipin.** *J Cell Biol* 2018.
Using cryo-EM, the authors report the structure of the luminal domain of *Drosophila* seipin, which exists as a ring-shaped dodecamer. In vitro experiments indicate that the luminal domain contains hydrophobic helices that can bind monolayers of LDs.
68. Ozeki S, Cheng J, Tauchi-Sato K, Hatano N, Taniguchi H, Fujimoto T: **Rab18 localizes to lipid droplets and induces their close apposition to the endoplasmic reticulum-derived membrane.** *J Cell Sci* 2005, **118**:2601-2611.
69. Pol A, Martin S, Fernandez MA, Ferguson C, Carozzi A, Luetterforst R, Enrich C, Parton RG: **Dynamic and regulated association of Caveolin with lipid bodies : modulation of lipid body motility and function by a dominant negative mutant.** *Mol Biol Cell* 2004, **15**:99-110.
70. Xu D, Li Y, Wu L, Li Y, Zhao D, Yu J, Huang T, Ferguson C, Parton RG, Yang H *et al.*: **Rab18 promotes lipid droplet (LD) growth by tethering the ER to LDs through SNARE and NRZ interactions.** *J Cell Biol* 2018, **217**:975-995.
The authors showed that Rab3GAP1/2, the GEF of Rab18, recruits and targets Rab18 to LDs. There, it interacts with the NAG-RINT1-ZW10 (NRZ) tethering complex and their associated SNAREs in the ER. This complex can bring the ER and LDs closer together. Depletion of Rab18 or associated components impaired neutral lipid storage and LD maturation.
71. Pulido MR, Diaz-Ruiz A, Jiménez-Gómez Y, Garcia-Navarro S, Gracia-Navarro F, Tinahones F, López-Miranda J, Frühbeck G, Vázquez-Martínez R, Malagón MM: **Rab18 dynamics in adipocytes in relation to lipogenesis, lipolysis and obesity.** *PLoS One* 2011, **6**:e22931.
72. Li C, Luo X, Zhao S, Siu GK, Liang Y, Chan HC, Satoh A, Yu SS: **COPI-TRAPP II activates Rab18 and regulates its lipid droplet association.** *EMBO J* 2017, **36**:441-457.
73. Jayson CBK, Arlt H, Fischer AW, Lai ZW, Farese RV, Walther TC: **Rab18 is not necessary for lipid droplet biogenesis or turnover in human mammary carcinoma cells.** *Mol Biol Cell* 2018, **29**:2037-2136.
74. Xu N, Zhang SO, Cole Ra, McKinney Sa, Guo F, Haas JT, Bobba S, Farese RV, Mak HY: **The FATP1-DGAT2 complex facilitates lipid droplet expansion at the ER-lipid droplet interface.** *J Cell Biol* 2012, **198**:895-911.
75. Weber-Boyvat M, Kentala H, Peränen J, Olkkonen VM: **Ligand-dependent localization and function of ORP-VAP complexes at membrane contact sites.** *Cell Mol Life Sci* 2015, **72**:1967-1987.
76. Kentala H, Koponen A, Vihinen H, Pirhonen J, Liebisch G, Pataj Z, Kivelä A, Li S, Karhinen L, Jääskeläinen E *et al.*: **OSBP-related protein-2 (ORP2): a novel Akt effector that controls cellular energy metabolism.** *Cell Mol Life Sci* 2018, **75**:1-17.
77. Tabara LC, Escalante R: **VMP1 establishes ER-microdomains that regulate membrane contact sites and autophagy.** *PLoS One* 2016, **11**:e0166499.
78. Markgraf DF, Klemm RW, Junker M, Hannibal-Bach HK, Ejsing CS, Rapoport TA: **An ER protein functionally couples neutral lipid metabolism on lipid droplets to membrane lipid synthesis in the ER.** *Cell Rep* 2014, **6**:44-55.